

# UNIVERSALITY OF COPRODUCTS IN CATEGORIES OF LAX ALGEBRAS

MOJGAN MAHMOUDI, CHRISTOPH SCHUBERT, AND WALTER THOLEN

ABSTRACT. Categories of lax  $(T, V)$ -algebras are shown to have pullback-stable coproducts if  $T$  preserves inverse images. The general result not only gives a common proof of this property in many topological categories but also shows that important topological categories, like the category of uniform spaces, are not presentable as a category of lax  $(T, V)$ -algebras, with  $T$  preserving inverse images. Moreover, we show that any such category of  $(T, V)$ -algebras has a concrete, coproduct preserving functor into the category of topological spaces.

Universality of coproducts is a property that distinguishes **Set**-based topological categories: while in many “everyday” topological categories coproducts are stable under pullback (topological spaces, preordered sets, premetric spaces, approach spaces), some others fail to enjoy the property (uniform spaces, proximity spaces, nearness spaces, merotopic spaces, see [6]). All of the topological categories in the first group happen to be presentable as categories of lax algebras, that is: they are of the form  $\mathbf{Alg}(T, V)$ , for a suitable extension  $T$  of a **Set**-monad  $T_0 = (T_0, e, m)$  and a complete lattice  $V$  that comes with an associative and commutative binary operation  $\otimes$  preserving suprema in each variable, and a  $\otimes$ -neutral element  $k$  (distinct from the bottom element  $\perp$ ), [5]. In this note, we show that this observation is not coincidental, that is: coproducts in  $\mathbf{Alg}(T, V)$  are always stable under pullback, making  $\mathbf{Alg}(T, V)$  in fact an (infinitely) extensive category, provided that  $T_0$  preserves inverse images. This condition is weaker than preservation of the Beck–Chevalley Property, as used by Clementino and Hofmann (see [4]); for an example, see [7, Example 1.3(a)].

Of special importance are *open morphisms* of lax algebras, as defined by Möbus [9] in the context of relational algebras in a category, and by Clementino and Hofmann [4] in the context used in this paper. These generalize open morphisms in the category **Top** of topological space. We characterize coproducts in  $\mathbf{Alg}(T, V)$  by the fact that all injections of the underlying **Set**-coproduct are open morphisms in  $\mathbf{Alg}(T, V)$ . Moreover, we construct a concrete functor  $\mathbf{Alg}(T, V) \rightarrow \mathbf{Top}$  which preserves open embeddings and, hence, coproducts. This result generalizes a construction of Manes [8].

---

*Date:* May 8, 2006.

*1991 Mathematics Subject Classification.* 18C99, 18A30.

*Key words and phrases.* Lax  $(T, V)$ -algebra, universality of coproducts, extensive category, topological category.

**Acknowledgments.** The first author thanks the Department of Mathematics and Statistics at York University for their kind hospitality and financial support during her stay in the Winter of 2005. She also acknowledges financial support of Shahid Beheshti University. The second author gratefully acknowledges a PhD-scholarship from the Ernst A.-C. Lange Stiftung, Bremen. The third author acknowledges financial support from NSERC.

## 1. PRELIMINARIES

Recall that for sets  $X, Y$ , a  $V$ -relation (or  $V$ -matrix)  $r : X \dashrightarrow Y$  from  $X$  to  $Y$  is a mapping  $r : X \times Y \rightarrow V$ ; it gets composed with  $s : Y \dashrightarrow Z$  via

$$(sr)(x, z) = \bigvee_{y \in Y} r(x, y) \otimes s(y, z).$$

The category  $\mathbf{Mat}(V)$  of sets and  $V$ -matrices is, using the pointwise order on the hom-sets, enriched over the monoidal closed category  $\mathbf{Sup}$  of complete lattices and suprema-preserving morphisms. That is, every hom-set  $\mathbf{Mat}(V)(X, Y)$  is a complete lattice, and the composition functions preserve suprema in each variable:

$$s(\bigvee r_i) = \bigvee sr_i, \quad (\bigvee r_i)q = \bigvee r_iq.$$

Every  $\mathbf{Set}$ -map  $f : X \rightarrow Y$  can be considered as a  $V$ -relation via  $f(x, y) = k$  if  $f(x) = y$ , and  $f(x, y) = \perp$  else. We also use the converse  $V$ -relation  $f^* : Y \dashrightarrow X$  with  $f^*(y, x) = f(x, y)$  and take note of the inequalities  $1_X \leq f^*f$ ,  $ff^* \leq 1_Y$ , which exhibit  $f^*$  as right adjoint to  $f$  in the 2-category  $\mathbf{Mat}(V)$ .

Let  $T_0 = (T_0, e, m)$  be a  $\mathbf{Set}$ -monad such that  $T_0$  preserves inverse images; that is:  $T_0$  preserves every  $\mathbf{Set}$ -pullback

$$\begin{array}{ccc} f^{-1}[B] & \longrightarrow & B \\ \downarrow & & \downarrow m \\ X & \xrightarrow{f} & Y \end{array}$$

when  $m$  is a monomorphism. In particular,  $T_0$  preserves monomorphisms.

Examples for inverse image-preserving monads abound; for instance, the powerset monad, the filter monad, the ultrafilter monad, and the free-monoid monad all preserve inverse images. But there are  $\mathbf{Set}$ -monads  $(T_0, e, m)$  for which  $T_0$  does not preserve inverse images: consider, for example, the trivial monad with  $T_0\emptyset = \emptyset$  and  $T_0X = 1$  for all  $X \neq \emptyset$ .

We now consider a lax extension  $T : \mathbf{Mat}(V) \rightarrow \mathbf{Mat}(V)$  of  $T_0$  to  $V$ -relations, which respects the order of  $V$ -relations and satisfies the following conditions

- (0)  $T_0f \leq Tf$  and  $(T_0f)^* \leq Tf^*$ ,
- (1)  $e_Y r \leq (Tr)e_X$ ,
- (2)  $m_Y(T^2r) \leq (Tr)m_X$ ,
- (3)  $(Ts)(Tr) \leq T(sr)$ ,

for all  $f : X \rightarrow Y$ ,  $r : X \rightrightarrows Y$ ,  $s : Y \rightrightarrows Z$ . On objects  $T$  coincides with  $T_0$ . As noted in [10], these conditions suffice to derive the identities

$$(4) \quad (Ts)(Tf) = T(sf) = (Ts)(T_0f),$$

$$(5) \quad (Tg^*)(Tr) = T(g^*r) = (T_0g)^*(Tr)$$

with  $g : Y \rightarrow Z$ , which play a crucial role in [5]. Moreover, we assume that

$$(6) \quad T(rg^*) = (Tr)(T_0g^*)$$

whenever  $g$  is a *monomorphism*.

## 2. GRAPHS AND LAX ALGEBRAS

The category  $\mathbf{Gph}(T, V)$  of  $(T, V)$ -graphs has as objects sets  $X$  equipped with a  $V$ -relation  $a : TX \rightrightarrows X$  satisfying  $1_X \leq ae_X$ ; a morphism  $f : (X, a) \rightarrow (Y, b)$  is a  $\mathbf{Set}$ -map  $f : X \rightarrow Y$  satisfying  $fa \leq bT_0f$ . The forgetful functor  $\mathbf{Gph}(T, V) \rightarrow \mathbf{Set}$  is topological: for a (possibly large) family  $(Y_i, b_i)_{i \in I}$  of  $(T, V)$ -graphs and mappings  $f_i : X \rightarrow Y_i$ , the so-called *initial structure*  $a$  on  $X$  (see [1]) is given by

$$a = \bigwedge_{i \in I} f_i^* b_i (T_0 f_i),$$

(see [5]). Hence  $\mathbf{Gph}(T, V)$  is complete and cocomplete. In particular,  $\mathbf{Gph}(T, V)$  has pullbacks, and they are formed by putting the initial structure with respect to the two projections on the  $\mathbf{Set}$ -pullback.

Colimits in  $\mathbf{Gph}(T, V)$  are formed as follows: given a diagram with vertices  $(X_i, a_i)_{i \in I}$ , first form the colimit  $(g_i : X_i \rightarrow Y)$  of the underlying diagram in  $\mathbf{Set}$ , and then equip  $Y$  with the *final structure*

$$(1) \quad b = \bigvee_{i \in I} g_i a_i (T_0 g_i)^*.$$

It is easy to see that  $1_Y \leq be_Y$  is satisfied by virtue of  $(g_i)_{i \in I}$  being jointly epic; equivalently, since  $\bigvee_I g_i g_i^* = 1_Y$ .

A  $(T, V)$ -graph  $(X, a)$  satisfies the inequality  $a \leq a(Ta)m_X^*$ ; it is called a *(lax)  $(T, V)$ -algebra* if it satisfies the transitivity condition  $a(Ta)m_X^* \leq a$  or, equivalently,  $a(Ta) \leq am_X$ . The full subcategory of  $\mathbf{Gph}(T, V)$  formed by the algebras is denoted by  $\mathbf{Alg}(T, V)$ . It is concretely reflective, with the reflector defined as follows (see [3]): given a graph  $(X, a)$ , form the transfinite sequence of  $V$ -relations

$$(2) \quad a_0 = a, \quad a_{\alpha+1} = a_\alpha(Ta_\alpha)m_X^*, \quad a_\gamma = \bigvee_{\beta < \gamma} a_\beta$$

for all ordinals  $\alpha$  and limit ordinals  $\gamma$ . One obtains an ascending chain of  $V$ -relations, which has to become stationary at some  $\alpha_0$ , we write  $L(X, a)$  for  $(X, a_{\alpha_0})$ . Then  $1_X : (X, a) \rightarrow L(X, a)$  is the sought universal arrow.

Thus also  $\mathbf{Alg}(T, V)$  is topological, limits in  $\mathbf{Alg}(T, V)$  are formed like in  $\mathbf{Gph}(T, V)$ , and colimits are formed by applying the reflector to the colimit in  $\mathbf{Gph}(T, V)$ .

## 3. COPRODUCTS

For any coproduct  $(t_i : X_i \rightarrow X)$  in **Set**, we easily obtain  $t_i^* t_i = 1_{X_i}$  and  $t_i^* t_j = \perp$  if  $i \neq j$  as well as  $\bigvee t_i^* t_i = 1_X$ . In fact, these equations make  $(t_i : X_i \rightarrow X)$  a *biproduct* in  $\mathbf{Mat} V$ .

Consequently, for any coproduct  $(t_i : (X_i, a_i) \rightarrow (X, c))$  of graphs,

$$(3) \quad t_i^* c = t_i^* \bigvee t_j a_j (T_0 t_j)^* = \bigvee t_i^* t_j a_j (T_0 t_j)^* = a_i (T_0 t_i)^*.$$

In other words, each injection  $t_i : (X_i, a_i) \rightarrow (X, c)$  is open in the sense of [4]: a morphism  $f : (X, a) \rightarrow (Y, b)$  is *open* if  $f^* b = a (T_0 f)^*$ . Moreover,  $t_i^* c T_0 t_i = a_i (T_0 t_i)^* T_0 t_i = a_i$  holds since  $T_0$  since preserves monomorphisms. Therefore, each open injection is an *embedding*, i.e. its domain carries the initial structure.

To study coproducts of  $(T, V)$ -algebras, we first note:

**Lemma 1.** *If  $f : (X, a) \rightarrow (Y, b)$  is an open injection from a  $(T, V)$ -algebra into a  $(T, V)$ -graph, then  $f : (X, a) \rightarrow L(Y, b)$  is also open.*

*Proof.* It suffices to show that  $f^* b_\alpha = a (T_0 f)^*$  holds for all ordinals  $\alpha$ . Indeed, the initial step is trivial, and for a successor ordinal we have:

$$\begin{aligned} f^* b_{\alpha+1} &= f^* b_\alpha (T b_\alpha) m_Y^* \\ &= a (T_0 f)^* (T b_\alpha) m_Y^* \\ &= a T (f^* b_\alpha) m_Y^* \\ &= a T (a (T_0 f)^*) m_Y^* \\ &= a (T a) (T_0 T_0 f)^* m_Y^* \\ &= a (T a) m_X^* (T_0 f)^* \\ &= a (T_0 f)^*. \end{aligned}$$

In case  $\beta$  is a limit ordinal, we have  $f^* b_\beta = f^* \bigvee_{\alpha < \beta} b_\alpha = \bigvee_{\alpha < \beta} f^* b_\alpha = \bigvee_{\alpha < \beta} a (T_0 f)^* = a (T_0 f)^*$ .  $\square$

**Theorem 2.** *Let  $(X_i, a_i)_I$  be a family of  $(T, V)$ -algebras and  $(t_i : X_i \rightarrow X)_I$  be a **Set**-coproduct. The following statements are equivalent for a  $(T, V)$ -graph  $(X, a)$ :*

- (1)  $(t_i : (X_i, a_i) \rightarrow (X, a))_I$  is a coproduct in  $\mathbf{Gph}(T, V)$ ;
- (2)  $(X, a) \in \mathbf{Alg}(T, V)$ , and  $(t_i : (X_i, a_i) \rightarrow (X, a))_I$  is a coproduct in  $\mathbf{Alg}(T, V)$ ;
- (3) each  $t_i : (X_i, a_i) \rightarrow (X, a)$  is open.

*Proof.* (1) implies (2): Let  $(X, c)$  denote the coproduct of the  $(X_i, a_i)$  in  $\mathbf{Alg}(T, V)$ . By Lemma 1, each  $t_i : (X_i, a_i) \rightarrow (X, c)$  is open. Thus  $c = \bigvee t_i^* t_i c = \bigvee t_i^* a_i (T_0 t_i)^*$ , and this is, by hypothesis, precisely the final structure  $a$  with respect to the coproduct injections.

(2) implies (3): Lemma 1.

(3) implies (1): From (3) one obtains  $a = \bigvee t_i t_i^* a = \bigvee t_i a_i (T_0 t_i)^*$ , so that, again,  $a$  is the final structure with respect to the coproduct injections.  $\square$

**Lemma 3.** *The following statements are equivalent for a set-indexed family  $(f_i : (X_i, a_i) \rightarrow (Y, b))_I$  of morphisms in  $\text{Alg}(T, V)$ :*

- (1) *each  $f_i$  is open;*
- (2) *the induced morphism  $f : \coprod_I (X_i, a_i) \rightarrow (Y, b)$  is open.*

*Proof.* (1) implies (2): Let  $(t_i : (X_i, a_i) \rightarrow (X, a))_I$  be a coproduct in  $\text{Alg}(T, V)$ . Using the easily verified formula  $f^* = \bigvee t_i f_i^*$ , we obtain:

$$a(T_0 f)^* = \bigvee t_i a_i (T_0 t_i)^* (T_0 f)^* = \bigvee t_i a_i (T_0 f_i)^* = \bigvee t_i f_i^* b = f^* b$$

and thus  $f$  is open.

(2) implies (1): Follows from the closedness of open morphisms under composition.  $\square$

In particular, open morphisms are closed under coproducts.

**Lemma 4.** *Openness of embeddings is stable under pullback in  $\text{Alg}(T, V)$ .*

For the proof we recall two well-known properties of  $\text{Mat } V$ :

**Remark 5.** *Set-pullbacks are Beck–Chevalley squares in  $\text{Mat } V$ . That is, given a set pullback*

$$\begin{array}{ccc} P & \xrightarrow{p} & Y \\ q \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

$f^* g = q p^*$  holds in  $\text{Mat } V$ .

*Proof.* Straightforward.  $\square$

**Remark 6.**  *$\text{Mat } V$  satisfies the following weak form of Freyd’s Modular Law: Given  $V$ -matrices  $r : X \rightarrow Z$ ,  $s : Y \rightarrow Z$ , and an injective function  $m : X \rightarrow Y$ , we have*

$$(4) \quad (r \wedge sm)m^* = rm^* \wedge s.$$

*Proof.* We assume  $m$  to be a subset-inclusion, and obtain:

$$rm^*(y, z) = \begin{cases} r(y, z) & \text{if } y \in X, \\ \perp & \text{else.} \end{cases}$$

Thus the left-hand side as well as the right-hand side of (4) evaluate to  $\perp$  for every  $(y, z)$  with  $y \notin X$ . Obviously both evaluate to  $r(y, z) \wedge s(y, z)$  for all  $(y, z)$  such that  $y \in X$ , and hence (4) holds.  $\square$

*Proof of Lemma 4.* Suppose

$$\begin{array}{ccc} (P, d) & \xrightarrow{p} & (Y, b) \\ q \downarrow & & \downarrow g \\ (X, a) & \xrightarrow{f} & (Z, c) \end{array}$$

is a pullback diagram in  $\mathbf{Alg}(T, V)$  and  $f : (X, a) \rightarrow (Z, c)$  is an open embedding. We have to show that also  $p : (P, d) \rightarrow (Y, b)$  is open. Observe that with  $f$  also  $T_0p$  is injective, and that the  $T_0$  image of the underlying Set-pullback is still a pullback, hence a Beck–Chevalley square. Thus we obtain  $q^*a(T_0q)(T_0p)^* = q^*a(T_0f)^*(T_0g) = q^*f^*c(T_0g) = p^*g^*c(T_0g)$ , hence

$$\begin{aligned} d(T_0p)^* &= (p^*b(T_0p) \wedge q^*a(T_0q))(T_0p)^* \\ &= p^*b \wedge q^*a(T_0q)(T_0p)^* \\ &= p^*b \wedge p^*g^*c(T_0g) \\ &= p^*(b \wedge g^*c(T_0g)) \\ &= p^*b. \end{aligned}$$

□

Lemma 4, together with Theorem 2 and the fact that coproducts in  $\mathbf{Set}$  are universal, immediately yields:

**Theorem 7.** *Coproducts in  $\mathbf{Alg}(T, V)$  are universal; that is: coproducts are stable under pullback.*

**Corollary 8.**  *$\mathbf{Alg}(T, V)$  is (infinitely) extensive.*

*Proof.* Coproducts in  $\mathbf{Alg}(T, V)$  are trivially disjoint, and this property in conjunction with universality gives extensivity (see [2]). □

Since coproducts fail to be universal in  $\mathbf{Unif}$  (see [6]), we also obtain:

**Corollary 9.** *The category of uniform spaces is not presentable in the form  $\mathbf{Alg}(T, V)$  with  $T_0$  preserving inverse images.*

Note, however, that  $\mathbf{Unif}$  is a coreflective subcategory of a category of lax  $(T, V)$ -proalgebras (see [5]).

#### 4. THE TOPOLOGY OF A $(T, V)$ -ALGEBRA

**Lemma 10.**  *$gf$  open and  $f$  surjective implies  $g$  open.*

*Proof.* Let  $f : (X, a) \rightarrow (Y, b)$  and  $g : (Y, b) \rightarrow (Z, c)$  be morphisms such that  $gf$  is open and  $f$  is surjective; equivalently,  $ff^* = 1_Y$ . It suffices to show  $g^*c \leq b(T_0g)^*$ :  $g^*c = ff^*g^*c = fa(T_0f)^*(T_0g)^*$  holds since  $gf$  is open, and thus, since  $f$  is a morphism,  $g^*c \leq b(T_0g)^*$ . □

For a  $(T, V)$ -algebra  $(X, a)$ , we write  $\mathcal{O}(X, a)$  for the set of all open subalgebras of  $(X, a)$ , that is,

$$\mathcal{O}(X, a) = \{ \iota_A : A \hookrightarrow X \mid \iota_A \text{ is open w.r.t. the initial structure} \}.$$

**Lemma 11.**  *$\mathcal{O}(X, a)$  is closed under arbitrary suprema and finite infima in the powerset  $PX$  of  $X$ .*

*Proof.* Suprema in  $PX$  can be described as follows: given a family  $(\iota_{A_i} : A_i \hookrightarrow X)_I$  of subset-inclusions, first form the induced morphism  $h = [\iota_{A_i}] : \coprod A_i \rightarrow X$ , and then factor  $h = \iota p$  with  $\iota$  a subset-inclusion and  $p$  surjective.  $\iota$  is the sought supremum.

Now if each  $\iota_{A_i}$  is open, so is  $h$  by Lemma 3, and thus  $\iota$  is open by Lemma 10.

By Lemma 4, the set  $\mathcal{O}(X, a)$  is closed under binary intersections, and since every isomorphism is open, this implies closedness under finite intersections.  $\square$

Hence  $\mathcal{O}(X, a)$  is a topology on  $X$ . Moreover, since open embeddings are stable under pullback in  $\text{Alg}(T, V)$ , every lax morphism  $f : (X, a) \rightarrow (Y, b)$  gives rise to a continuous function  $f : \mathcal{O}(X, a) \rightarrow \mathcal{O}(Y, b)$ . Thus:

**Theorem 12.** *There exists a concrete functor  $O : \text{Alg}(T, V) \rightarrow \text{Top}$ , defined by*

$$(X, a) \mapsto (X, \mathcal{O}(X, a)).$$

*$O$  preserves open embeddings and coproducts.*

*Proof.* Clearly  $O$  is a concrete functor which preserves open embeddings. Preservation of coproducts follows thus from the characterization of coproducts in  $\text{Alg}(T, V)$  (and hence in  $\text{Top}$ ) given in Theorem 2.  $\square$

## REFERENCES

- [1] Adamek, J., H. Herrlich and G. Strecker. *Abstract and Concrete Categories*, John Wiley and Sons, New York, 1990. (Available on-line at <http://katmat.math.uni-bremen.de/acc/acc.pdf>).
- [2] Carboni, A., S. Lack, R. F. C. Walters. Introduction to extensive and distributive categories, *J. Pure Appl. Algebra* **84** (1993), 145–158.
- [3] Clementino, M. M., D. Hofmann. Topological features of lax algebras, *Appl. Categ. Structures* **11** (2003), 267–286.
- [4] Clementino, M. M., D. Hofmann. Effective descent morphisms in categories of lax algebras, *Appl. Categ. Structures* **12** (2004), 413–425.
- [5] Clementino, M. M., D. Hofmann, W. Tholen. One setting for all: Metric, Topology, Uniformity, Approach Structure, *Appl. Categ. Structures* **12** (2004), 127–154.
- [6] Herrlich, H. Are there convenient subcategories of TOP?, *Topology Appl.* **15** (1983), 263–271
- [7] Johnstone, P., J. Power, T. Tsujishita, H. Watanabe, J. Worrell. On the structure of categories of coalgebras. *Theor. Comp. Science* **260** (2001), 87–117.
- [8] Manes, E. Taut Monads and  $T0$ -spaces, *Theor. Comp. Science* **275** (2002), 79–109.
- [9] Möbus, A. Relational-Algebren, Ph.D. thesis (University of Düsseldorf, 1981).

- [10] Seal, G. J. Canonical and op-canonical lax algebras, *Theory Appl. Categories* **14** (2005), 221–243.

DEPARTMENT OF MATHEMATICS, SHAHID BEHESHTI UNIVERSITY, TEHRAN 19839,  
IRAN

*E-mail address:* `m-mahmoudi@sbu.ac.ir`

SOFTWARE-TECHNOLOGY, FACHBEREICH INFORMATIK, UNIVERSITÄT DORTMUND, 44221  
DORTMUND, GERMANY

*E-mail address:* `christoph.schubert@uni-dortmund.de`

*URL:* `http://ls10-www.cs.uni-dortmund.de/~schubert`

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ON  
M3J 1P3, CANADA

*E-mail address:* `tholen@mathstat.yorku.ca`